

## **Nerve Blocks**

# **READING LIST**

#### Chapter 1

Brull, R, McCartney, CJ, Chan, VW, et al. 2007. Neurological complications after regional anesthesia: contemporary estimates of risk. *Anesth Analg.* **104**: 965–974.

Cass, SP. 2016. Ultrasound-guided nerve hydrodissection: what is it? A review of the literature. *Curr Sports Med Rep.* **15**: 20–22.

Fredrickson MJ, and Kilfoyle DH. 2009. Neurologic complication analysis of 1000 ultrasound guided peripheral nerve blocks for elective orthopedic surgery: a prospective study. *Anesthesia*. **64**: 836–844.

Liu, SS, YaDeau, JT, Shaw, PM, et al. 2011. Incidence of unintentional intraneural injection and postoperative neurological complications with ultrasound-guided interscalene and supraclavicular nerve blocks. *Anaesthesia*. 66: 168–174.

Marhofer, P, Greher, M, and Kapral, S. 2004. Ultrasound guidance in regional anaesthesia. *Br J Anaesth*. **94**: 7–17.

Martinoli, C, Bianchi, S, Dahmane, MH, et al. 2002. Ultrasound of tendons and nerves. Eur Radiol. 12: 44-55.

Walker, FO, Cartwright, MS, Wiesler, ER, et al. 2004. Ultrasound of nerve and muscle. *Clin Neurophysiol.* **115**: 495–507.

#### **Chapter 2**

Arcand, G, Williams, SR, Chouinard, P, et al. 2005. Ultrasound-guided infraclavicular versus supraclavicular block. *Anesth Analg.* **101**: 886–890.

Chan, VW, Perlas, A, Rawson, R, et al. 2003. Ultrasound-guided supraclavicular brachial plexus block. *Anesth Analg.* **97**: 1514–1517.

Fredrickson MJ, Ball CM, and Dalgleish AJ. 2011. Posterior versus Anterolateral Approach Interscalene Catheter Placement: A Prospective Randomized Trial. *Reg Anesth Pain Med.* 36: 125–133.

Kapral, S, Krafft, P, Eibenberger, K, et al. 1994. Ultrasound-guided supraclavicular approach for regional anesthesia of the brachial plexus. *Anesth Analg.* **78**: 507–513.

Riazi, S, Carmichael, N, Awad, I, et al. 2008. Effect of local anaesthetic volume (20 vs 5 ml) on the efficacy and respiratory consequences of ultrasound-guided interscalene brachial plexus block. *Br J Anaesth*. **101**: 549–556.

Sandhu, NS and Capan, LM. 2002. Ultrasound-guided infraclavicular brachial plexus block. *Br J Anaesth*. **89**: 254–259.

Spence, BC, Beach, ML, Gallagher, JD, et al. 2011. Ultrasound-guided interscalene blocks: understanding where to inject the local anaesthetic. *Anaesthesia*. **66**: 509–514.



Williams, SR, Chouinard, P, Arcand, G, et al. 2003. Ultrasound guidance speeds execution and improves the quality of supraclavicular block. *Anesth Analg.* **97**: 1518–1523.

#### **Chapter 3**

Baaj, JM, Alsatli, RA, Majaj, HA, et al. 2010. Efficacy of ultrasound-guided transversus abdominis plane (TAP) block for postcesarean section delivery analgesia--a double-blind, placebo-controlled, randomized study. *Middle East J Anaesthesiol.* **20**: 821–826.

Bergeron E, Lavoie A, Clas D, et al. 2003. Elderly trauma patients with rib fractures are at greater risk of death and pneumonia. *J Trauma*. **54**: 478–485.

Blanco, R, Parras, T, McDonnell, JG, et al. 2013. Serratus plane block: a novel ultrasound-guided thoracic wall nerve block. *Anesthesia*. **68**:1107–1113.

Durant, E, Dixon, B, Luftig, J, et al. 2017. Ultrasound-guided serratus plane block for ED rib fracture pain control. *Am J Emerg Med.* **35**: 197.e3–197.e6.

Hebbard, P, Fujiwara, Y, Shibata, Y, et al. 2007. Ultrasound-guided transversus abdominis plane (TAP) block. *Anaesth Intensive Care*. **35**: 616–618.

McDonnell, JG, Curley, G, Carney, J, et al. 2008. The analgesic efficacy of transversus abdominis plane block after cesarean delivery: a randomized controlled trial. *Anesth Analg.* **106**: 186–191.

Peterson, MK, Millar, FA, and Sheppard, DG, 2002. Editorial I: Ultrasound-guided nerve blocks. *Br J Anaesth*. **88**: 621–624.

Walter, CJ, Maxwell-Armstrong, C, Pinkney, TD, et al. 2013. A randomised controlled trial of the efficacy of ultrasound-guided transversus abdominis plane (TAP) block in laparoscopic colorectal surgery. *Surg Endosc.* 27: 2366–2372.

#### Chapter 4

Blaivas, M, Adhikari, S, and Lander, L. 2011. A prospective comparison of procedural sedation and ultrasound guided interscalene nerve block for shoulder reduction in the emergency department. *Acad Emerg Med.* 18: 922–927.

Blaivas, M and Lyon, M. 2006. Ultrasound-guided interscalene block for shoulder dislocation reduction in the ED. *Am J Emerg Med.* **24**: 293–296.

Blakeley, C, Spencer, O, Newman-Saunders, T, et at. 2009. A novel use of portable ultrasound in the management of shoulder dislocation. *Emerg Med J.* **26**: 662–663.

Foxall, GL, Skinner, D, Hardman, JG, et al. 2007. Ultrasound anatomy of the radial nerve in the distal upper arm. *Reg Anesth Pain Med.* **32**: 217–220.

Gray, AT and Schafhalter-Zoppoth, I. 2003. Ultrasound guidance for ulnar nerve block in the forearm. *Reg Anesth Pain Med*. **28**: 335-339.

Liebmann, O, Price, D, Mills, C, et al. 2006. Feasibility of forearm ultrasonography-guided nerve blocks of the radial, ulnar, and median nerves for hand procedures in the emergency department. *Ann Emerg Med.* **48**: 558–562.

Ponrouch, M, Bouic, N, Bringuier, S, et al. 2010. Estimation and pharmacodynamic consequences of the minimum effective anesthetic volumes for median and ulnar nerve blocks: a randomized, double-blind, controlled comparison between ultrasound and nerve stimulation guidance. *Anesth Analg.* 111: 1059–1064.



### **Chapter 5**

Foss, NB, Kristensen, BB, Bundgaard, M, et al. 2007. Fascia iliaca compartment blockade for acute pain control in hip fracture patients: a randomized, placebo-controlled trial. *Anesthesiology*. **106**: 773–778.

Groot, L, Dijksman, LM, Simons, MP, et al. 2015. Single fascia iliaca compartment block is safe and effective for emergency pain relief in hip-fracture patients. *West J Emerg Med.* 16: 1188–1193.

Hanson, NA, Allen, CJ, Hostetter, LS, et al. 2014. Continuous ultrasound-guided adductor canal block for total knee arthroplasty: a randomized, double-blind trial. *Anesth Analg.* 118: 1370–1377.

Manickam, B, Perlas, A, Duggan, E, et al. 2009. Feasibility and efficacy of ultrasound-guided block of the saphenous nerve in the adductor canal. *Reg Anesth Pain Med*. **34**: 578–580.

Mouzopoulos, G, Vasiliadis, G, Lasanianos, N, et al. 2009. Fascia iliaca block prophylaxis for hip fracture patients at risk for delirium: a randomized placebo-controlled study. *J Orthop Traumatol.* **10**: 127–133.

Perlas, A, Brull, R, Chan, VW, et al. 2008. Ultrasound guidance improves the success of sciatic nerve block at the popliteal fossa. *Reg Anesth Pain Med.* 33: 259–265.

Reavley, P, Montgomery, AA, Smith, JE, et al. 2015. Randomised trial of the fascia iliaca block versus the '3-in-1' block for femoral neck fractures in the emergency department. *Emerg Med J.* **32**: 685–689.

Redborg, KE, Antonakakis, JG, Beach, ML, et al. 2009. Ultrasound improves the success rate of a tibial nerve block at the ankle. *Reg Anesth Pain Med.* **34**: 256–260.

Swenson, JD, Davis, JJ, Stream, JO, et al. 2015. Local anesthetic injection deep to the fascia iliaca at the level of the inguinal ligament: the pattern of distribution and effects on the obturator nerve. *J Clin Anesth.* 27: 652–657.

Wolfe, JW and Butterworth, JF. 2011. Local anesthetic systemic toxicity: update on mechanisms and treatment. *Curr Opin Anaesthesiol*. **24**: 561–566.

#### Chapter 6

Godwin, SA, Caro, DA, Wolf, SJ, et al. 2005. Clinical policy: procedural sedation and analgesia in the emergency department. *Ann Emerg Med.* **45**: 177–196.

Mariano, ER, Ilfeld, BM, Cheng, GS, et al. 2008. Feasibility of ultrasound-guided peripheral nerve block catheters for pain control on pediatric medical missions in developing countries. *Pediatr Anaesth.* 18: 598–601.

Oberndorfer, U, Marhofer, P, Bösenberg, A, et al. 2007. Ultrasonographic guidance for sciatic and femoral nerve blocks in children. *Br J Anaesth*. **98**: 797–801.

Paut, O, Sallabery, M, Schreiber-Deturmeny, E, et al. 2001. Continuous fascia iliaca compartment block in children: a prospective evaluation of plasma bupivacaine concentrations, pain scores, and side effects. *Anesth Analg.* **92**: 1159–1163.

Peña, BM and Krauss, B. 1999. Adverse events of procedural sedation and analgesia in a pediatric emergency department. *Ann Emerg Med.* **34**: 483–491.